

An ansatz for spacetimes of zero gravitational mass : global monopoles and textures

Naresh Dadhich*

*Inter-University Centre for Astronomy & Astrophysics,
Post Bag 4, Ganeshkhind, Pune - 411 007, India.*

*K. Narayan[†]Department of Physics,
Indian Institute of Technology Bombay,
Mumbai 400 076.*

Abstract

We propose a geometric ansatz, a restriction on Euclidean / Minkowski distance in the embedding space being proportional to distance in the embedded space, to generate spacetimes with vanishing gravitational mass ($R_{ik}u^i u^k = 0, u_i u^i = 1$). It turns out that these spacetimes can represent global monopoles and textures. Thus the ansatz is a prescription to generate zero mass spacetimes that could describe topological defects, global monopoles and textures.

PACS numbers: 04.20,04.60,98.80Hw

IUCAA-32/97

*E-mail : nkd@iucaa.ernet.in

[†]E-mail : narayan@niharika.phy.iitb.ernet.in

By the restriction that Euclidean / Minkowski distance in the embedding space is proportional to the distance in the embedded space we obtain spacetimes with the vanishing gravitational charge density defined by, $-4\pi\rho_c = R_{ik}u^i u^k = -4\pi(\rho + 3p)$, $u_i u^i = 1$. That means active gravitational mass of the spacetime is zero. This is also the characteristic property of spacetimes describing topological defects like global monopoles and global textures [1-4]. Such exotic objects are supposed to be created by phase transition in the early Universe when global symmetry is spontaneously broken. In recent times there have been some interesting applications of such objects in the early Universe cosmology [1-11].

Newtonian theory (NT) is a linear scalar theory, in which absence of gravitational field is characterised by constancy of the scalar potential, ϕ . The physical measurable effects of gravity are particle and tidal accelerations. For the canonical situation of field due to a mass point, the former is always radial while the latter is also experienced only by radially falling nearby particles. They are given by first and second derivatives of potential relative to radial coordinate r and hence both vanish when $\phi = \text{const}$.

Let us consider this question in general relativity (GR). How does one define potential in GR ? In the context of the Schwarzschild field, we would like to make the following working definition. By potential we shall mean a scalar function that (i) completely describes the field, (ii) is a solution of the Laplace equation corresponding to $R_t^t = 0$, (iii) defines equivalent Newtonian potential and (iv) determines radial acceleration for a free particle. Since GR is a non-linear theory, this scalar function also appears in the Riemann curvature as it is, which will in contrast to NT impart a physical meaning to it. This is the new and unique feature of GR, which does not consonant with the usual concept of potential in classical physics. We shall in what follows mean by it as stated above, essentially a function that completely characterises the Schwarzschild field and shall call it relativistic potential.

In GR, gravitation is described by curvature of spacetime. Relativistic potential not only gives rise to the acceleration but it also “curves” the space part of the metric. Note that we could have acceleration non-zero with space part of the metric being flat. It can be shown that non-linear feature of gravitation (field being its own source) goes into curving the space part of

the metric, which survives even when potential is constant but not zero [12-14]. This is how constant potential in GR gives rise to curved spacetime. Of course this will have zero gravitational mass and free of radial acceleration as well as tidal acceleration for radial motion. The curvature shows up only in tidal acceleration for transverse motion, which was however absent in NT because space was flat. This spacetime was first considered by the author [15] as an example of a spacetime, having of 20 curvatures all but one non-zero. It gave rise to a stress-system $T_t^t = T_r^r \sim 1/r^2$, and the rest of stresses being zero. This did not accord to any conventional matter fields and hence was not pursued any further and forgotten. On the other hand this is precisely the stresses required to describe a global monopole [1]. It is remarkable that the spacetime arising from a constant relativistic potential can actually describe an exotic object like global monopole, which is supposed to be created by spontaneously breaking of global symmetry $O(3)$ into $U(1)$.

The simplest model of a global monopole is described by the Lagrangian [1],

$$L = \frac{1}{2} \partial_i \psi^a \partial^i \psi^a - \frac{\lambda}{4} (\psi^a \psi^a - \eta^2)^2 \quad (1)$$

where ψ^a is a triplet of scalar fields, $\psi^a = \eta f(r) x^a / r$. It has global $O(3)$ symmetry which is spontaneously broken to $U(1)$. Outside the monopole core $f = 1$, and ψ^a takes the vacuum value $\eta x^a / r$ and stresses will be given by

$$T_t^t = T_r^r = \frac{\eta^2}{r^2} \quad (2)$$

and rest of $T_i^k = 0$.

Let us consider the spherically symmetric metric,

$$ds^2 = A dt^2 - B dr^2 - r^2 d\Omega^2, d\Omega^2 = d\theta^2 + \sin^2 \theta d\varphi^2. \quad (3)$$

Now $T_{tr} = 0$ and $T_t^t = T_r^r$ lead to $A = B^{-1} = 1 + 2\phi(r)$ and the two equations

$$R_t^t = -\nabla^2 \phi = 0 \quad (4)$$

$$R_\theta^\theta = R_\varphi^\varphi = -\frac{2}{r^2} (r\phi)' \quad (5)$$

where a dash denotes derivative w.r.t. r .

The former is the good old Laplace equation admitting the well-known general solution $\phi = K - M/r$ while the latter determines the stresses (2) when $K = -4\pi\eta^2 \neq 0$. Thus the Schwarzschild black hole with global monopole charge is described by the metric [1],

$$ds^2 = (1 - 8\pi\eta^2 - \frac{2M}{r})dt^2 - (1 - 8\pi\eta^2 - \frac{2M}{r})^{-1}dr^2 - r^2d\Omega^2. \quad (6)$$

Note that $\phi = K \neq 0$ gives rise to curved spacetime representing stresses corresponding to a global monopole. It goes over to the Schwarzschild field when $K = 0$.

In this note we shall propose an ansatz to obtain spacetimes with vanishing gravitational charge density (zero mass). Let us begin with a 5-dimensional flat spacetime

$$ds^2 = dt^2 - dx_1^2 - dx_2^2 - dx_3^2 - dx_4^2 \quad (7)$$

and impose the restriction

$$x_1^2 + x_2^2 + x_3^2 + x_4^2 = k^2(x_1^2 + x_2^2 + x_3^2) \quad (8)$$

where k is a constant. Now elimination of x_4 will lead to the global monopole metric,

$$ds^2 = dt^2 - k^2dr^2 - r^2d\Omega^2 \quad (9)$$

which is equivalent to (6) with $k^2 = (1 - 8\pi\eta^2)^{-1}$ and $M = 0$. Here t has not participated in the restriction (8).

If we had instead considered the restriction

$$x_2^2 + x_3^2 + x_4^2 = k^2(x_2^2 + x_3^2) \quad (10)$$

which would have led to the metric for a cosmic string with deficit angle and vanishing Riemann curvature.

If we let t also in and consider

$$t^2 - x_1^2 - x_4^2 = k^2(t^2 - x_1^2) \quad (11)$$

which would lead to the metric

$$ds^2 = k^2 d\mu^2 - \mu^2 d\theta^2 - dx_2^2 - dx_3^2 \quad (12)$$

where $\mu^2 = t^2 - x_1^2$. This is also a flat spacetime with a deficit angle. It may be considered as Lorentzian version of cosmic string for $\mu^2 = t^2 - x_1^2$. In general a metric with a deficit angle can be obtained by letting any two of the variables participate in the ansatz.

Analogous to Lorentzian cosmic string, let us consider Lorentzian monopole by

$$t^2 - x_1^2 - x_2^2 - x_4^2 = k^2(t^2 - x_1^2 - x_2^2) \quad (13)$$

which will yield the metric,

$$ds^2 = k^2 d\mu^2 - \mu^2 d\theta^2 - \mu^2 \sin^2 \theta d\varphi^2 - dx_3^2 \quad (14)$$

where $\mu^2 = t^2 - x_1^2 - x_2^2$. It gives rise to

$$T_\mu^\mu = T_{x_3}^{x_3} = \frac{1 - k^2}{8\pi k^2 \mu^2}, R_{\theta\varphi}^{\theta\varphi} = \frac{1 - k^2}{k^2 \mu^2} \quad (15)$$

and other components vanishing.

This is similar to the monopole case. These stresses can be formally generated by the same Lagrangian (1) with r being replaced by μ and $T_\mu^\mu = T_{x_3}^{x_3} = \eta^2/\mu^2$. This can be looked upon as Lorentzian version of global monopole.

Finally if we let all the variables to participate in the ansatz,

$$t^2 - x_1^2 - x_2^2 - x_3^2 - x_4^2 = k^2(t^2 - x_1^2 - x_2^2 - x_3^2) \quad (16)$$

which will give the metric

$$ds^2 = k^2 d\mu^2 - \mu^2 d\chi^2 - \mu^2 \sin^2 \chi d\Omega^2 \quad (17)$$

with $\mu^2 = t^2 - x_1^2 - x_2^2 - x_3^2$. This will give rise to perfect fluid with the equation of state $\rho + 3p = 0$,

$$8\pi\rho = \frac{3(1-k^2)}{k^2\mu^2}, R^{\theta\varphi}_{\theta\varphi} = R^{\theta\chi}_{\theta\chi} = R^{\varphi\chi}_{\varphi\chi} = \frac{1-k^2}{k^2\mu^2}. \quad (18)$$

This will correspond to a spacetime of global texture [2,10]. Again the above stresses can formally be generated from the same Lagrangian (1) with r being replaced by the Lorentzian μ and we get

$$T^\mu_\mu = \frac{3\eta^2}{2\mu^2}, T^\chi_\chi = T^\theta_\theta = T^\varphi_\varphi = \frac{\eta^2}{2\mu^2}. \quad (19)$$

This is the equation of state $\rho + 3p = 0$ employed to characterise vacuum energy or a new kind of exotic matter called k-matter[5- 8]. Several authors have considered cosmological implications of decaying vacuum energy and time varying cosmological term[5-8, 16- 17, and others]. Here we have given an interesting and novel way of generating this kind of exotic vacuum energy distribution, characterised by the equation of state $\rho + 3p = 0$. This indicates that spacetime has zero active gravitational mass.

Thus we have given an elegant geometric ansatz for generation of spacetimes of global monopoles and textures with vanishing gravitational mass. The ansatz is simply the restriction, that “distance” in the embedding space is propotional to “distance” in the embedded space. This generates curvature in embedded spacetime corresponding to vanishing gravitational mass. This is also the characteristic property of topological defects, global monopoles and textures. Here we do formally obtain Lorentzian versions of cosmic strings and global monopoles. Such objects have not been considered in literature. Their physical interpretation is an open issue. They formally follow from the ansatz when Euclidean distance is replaced by Lorentzian distance in the field configuration. Global texture is on the other hand is a 4- dimentional generalization of global monopole. It can be considered as generalization of Lorentzian monopole as well.

It is obvious from the ansatz that all these spacetimes will be by definition embeddable in 5-dimentional Minkowski spacetime (i.e. they are of class I). What is perhaps not so obvious is the fact that they are not conformally flat.

It would be interesting to consider spacetimes conformal to these zero mass spacetimes and examine whether they represent any specific properties. One of us (ND) has studied spacetimes conformal to global monopole metric (9) and has shown that it picks up uniquely isothermal (density falling off as inverse square of the area radius with a linear equation of state) character of fluid sphere without boundary. It is a general result that a necessary and sufficient condition for an isothermal fluid sphere without boundary is that its metric is conformal to the monopole metric (9) [18-19]. We shall consider separately spacetimes conformal to Lorentzian global monopole and texture spacetimes [20].

Global texture metric (17) is clearly not conformally flat instead it is conformal to hyperbolic static (Einstein) metric with negative curvature. There do however exist conformally flat spacetime with the equation of state $\rho + 3p = 0$. It is given by

$$ds^2 = e^{2\alpha t}(dt^2 - dr^2 - r^2 d\Omega^2), \alpha = \text{const.} \quad (20)$$

while the global texture will have the form

$$ds^2 = e^{2\eta/k}(d\eta^2 - \frac{dR^2}{1 + R^2} - R^2 d\Omega^2). \quad (21)$$

The two are different spacetimes because the base metric for (20) is flat while for (21) it is hyperbolic Einstein spacetime. In the latter case, both base as well as conformal spacetimes obey the equation of state $\rho + 3p = 0$.

Since gravitational mass $R_{ik}u^i u^k = 0$ is zero, the relativistic potential is constant, making acceleration as well as tidal acceleration for radial motion vanish for these spacetimes, they should in some sense be looked upon as “minimally” curved. They are free of Newtonian gravity. It is interesting to see their association with spacetimes describing global monopoles and textures. The ansatz could be considered as a prescription to generate “minimally” curved spacetimes that describe topological defects, global monopoles and textures.

Acknowledgement : KN thanks Jawaharlal Nehru Centre for Advanced Research for an award of summer fellowship that facilitated this work and IUCAA for hospitality. ND thanks Ramesh Tikekar for a useful discussion on embedding.

References

- [1] M Barriola and A Vilenkin (1989) Phys. Rev. Lett. **63**, 341.
- [2] R L Davis (1987) Phys. Rev. **35**, 3705.
- [3] N Turok (1989) Phys. Rev. Lett. **63**, 2625.
- [4] D Harari and C Lousto (1990) Phys. Rev. **D42**, 2626.
- [5] A Vilenkin (1984) Phys. Rev. Lett. **53**, 1016.
- [6] R J Gott III and M J Rees (1987) Mon. Not. R. Astr. Soc. **227**, 453.
- [7] E W Kobl (1989) Ap.J. **344**, 543.
- [8] M Kamionkowski and N Toumbas (1996) Phys. Rev. Lett. **77**, 587.
- [9] V Sahni (1991) Phys. Rev. **D43**, R301.
- [10] D Nötzold (1991) Phys. Rev. **D43**, R961.
- [11] M V John and K B Joseph (1996) Phys. Lett. **B 87**, 466.
- [12] N Dadhich (1995) GR-14 Abstracts, A.98.
- [13] ——— (1997) On the Schwarzschild field, to be submitted.
- [14] N Dadhich, K Narayan and U Yajnik (1997), Schwarzschild black hole with global monopole charge, submitted.
- [15] N Dadhich (1970) Ph.D. Thesis, Pune University, unpublished.
- [16] W Chen and Y Wu (1990) Phys. Rev. **D41**, 695.
- [17] M V John and K B Joseph (1997) A low matter density decaying vacuum cosmology from complex metric, submitted.
- [18] N Dadhich (1996) Gen. Relativ. Grav. **28**, 1455.
- [19] ——— (1997) Isothermal fluid sphere : Uniqueness and conformal mapping, submitted.
- [20] N Dadhich and K Narayan (1997) under preparation.